

Effect of Fish Oil Addition to the Skim Milk-egg Yolk Extender on the Quality of Frozen-thawed Bali Bull Spermatozoa

by Aam Gunawan

Submission date: 28-May-2023 02:47PM (UTC+0500)

Submission ID: 2103543833

File name: 3_Effect_of_Fish_Oil_Addition_to_the_skim_milk_yolk.pdf (1.04M)

Word count: 3382

Character count: 18075

Effect of Fish Oil Addition to the Skim Milk-egg Yolk Extender on the Quality of Frozen-thawed Bali Bull Spermatozoa^[1]

Abdul MALIK¹✉ Muhammad SYARIFDJAYA¹ Aam GUNAWAN¹
Siti ERLINA¹ Achmad JAELANI¹ Dadang Budi WIBOWO²

^[1] This study was supported by funding routine from Islamic University of Kalimantan

¹ Department of Animal Science, Faculty of Agriculture, Islamic University of Kalimantan, Banjarmasin, South Kalimantan, INDONESIA

² Officer of Animal husbandry, District of Balangan, Province South Kalimantan, INDONESIA

Article Code: KVFD-2017-17556 Received: 14.02.2017 Accepted: 14.04.2017 Published Online: 19.04.2017

Citation of This Article

Malik A, Syarifdjaya M, Gunawan A, Erlina S, Jaelani A, Wibowo DB: Effect of fish oil addition to the skim milk-egg yolk extender on the quality of frozen-thawed Bali bull spermatozoa. *Kafkas Univ Vet Fak Derg*, 23 (4): 651-654, 2017. DOI: 10.9775/kvfd.2017.17556

Abstract

This study was conducted to investigate the effect of fish oil addition to the semen extender on post-thaw quality of Bali bull spermatozoa. Fish oil was added at the doses of 0 (control), 50, 100, 150 and 200 mg/100 mL to the skim milk-egg yolk extender. Supplementation of fish oil at the dose of 100 mg/mL significantly (P<0.05) increased both live sperm rate and motility, but its dose of 200 mg/mL caused a significant (P<0.05) decrease in the live sperm rate when compared with the control group. A significant (P<0.05) increment was observed in abnormality rate of frozen-thawed spermatozoa in comparison with the control group when all doses of fish oil except 50 mg/100 mL were added. In conclusion, addition of 100 mg fish oil to 100 mL of skim milk-egg yolk extender could be beneficial for the improvement of the quality of Bali bull spermatozoa after freeze-thawing.

Keywords: Bull semen, Fish oil, Cryopreservation, Live sperm, Motility

Süt Tozu-Yumurta Sarısı Sulandırıcısına Balık Yağı İlavesinin Dondurulmuş-Çözdürülmüş Bali Boğa Spermatozoonlarının Kalitesi Üzerine Etkisi

Özet

Bu çalışma sperma sulandırıcısına katılan balık yağının Bali boğa spermatozoonlarının çözündürme sonrası kalitesi üzerine etkisini araştırmak amacıyla yapıldı. Süt tozu-yumurta sarısı sulandırıcısına 0 (kontrol), 50, 100, 150 ve 200 mg/100 mL dozunda balık yağı ilave edildi. 100 mg/100 mL balık yağı ilavesi kontrol grubu ile karşılaştırıldığında hem canlı sperm oranını hem de motiliteyi önemli derecede (P<0.05) artırdı, ancak 200 mg/100 mL'lik doz canlı sperm oranında önemli (P<0.05) bir azalmaya neden oldu. 50 mg/100 mL'lik doz hariç balık yağının tüm dozları ilave edildiğinde kontrol grubuna kıyasla dondurulmuş-çözdürülmüş spermatozoonların anormallik oranında önemli (P<0.05) bir artış gözlemlendi. Sonuç olarak, süt tozu-yumurta sarısı sulandırıcısının 100 mL'sine 100 mg balık yağı ilavesi Bali boğa spermatozoonlarının dondurma-çözündürme sonrası kalitesinin artırılması için faydalı olabilir.

Anahtar sözcükler: Boğa sperması, Balık yağı, Kriyoprezervasyon, Canlı sperm, Motilite

INTRODUCTION

Artificial insemination (AI) is an important tool for genetic improvement and has effectively been used in several livestock species including cattle, sheep and goats. One of the factors affecting the success of AI is the semen cryopreservation. The good post-thawed semen quality depends on the composition of the cryopreservation media. The process of semen cryopreservation is a stage of procedure that causes several forms of spermatozoa

damages. One of the most significant components in semen processing is the ingredients of extender that provide protection to the sperm against cold shock^[1]. The injuries in post-thawed spermatozoa are generally associated with osmotic changes, membrane alteration, inter-and intra-cellular^[2,3] crystal formation during cryopreservation process^[2,3]. Therefore, cryoprotectants are supplemented to the semen extender to reduce damaging effects of cryopreservation processes^[4]. Considerable evidences suggest that lipidic composition of the sperm membrane is



İletişim (Correspondence)



+62 813 35213112



sidol_99@yahoo.com

responsible for the detrimental effect of cryopreservation¹⁵. Before semen freezing, percent docosahexaenoic acid was greater in fatty acid treatment than that in the group without fatty acid and reduced significantly in both groups after thawing¹⁶.

Fish oil contains omega 3 fatty acid and is a major source of eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA)¹⁴. However, it has been reported¹⁷ that fish oil contains 1970 mg saturated fatty acid (SFA), 35.5% polyunsaturated fatty acid (PUFA), 180 mg eicosapentaenoic acid (EPA) and 120 mg docosahexaenoic acid (DHA). Many researchers have assessed the effect of fish oil addition on semen quality parameters in domestic animals. They have reported that fish oil caused increments in sperm motility and reductions in abnormal sperm in boars¹⁸, improved fertility in turkeys¹⁹ as well as increased total of sperm count in rams²⁰. On the other hand, the addition of fish oil as a source of omega-3 fatty acids to diet of boars has been demonstrated to possess no significant improvement effect on the quality of frozen-thawed sperm²¹. Similarly, incorporation of DHA as an omega-3 fatty acid in the diet has been found to improve significantly the motility, live sperm²² and normal morphology of fresh semen^{23,24}. Addition of DHA from fish oil to the egg yolk extender²⁵ was found to be effective for increase the progressive motility and membrane integrity in post-thawed boar semen²⁶. Although there are many evidences about the influence of DHA enriched extenders on freezing of boar semen²⁷, however, there are limited numbers of reports about the effect of fish oil and/or DHA enriched extender on the quality of frozen-thawed bull semen^{28,29}. The objective of this study was to evaluate the effect of fish oil addition to the extender on live sperm rate, motility and abnormality of post-thawed Bali bull spermatozoa.

MATERIAL and METHODS

Semen Collection

Semen was collected by artificial vagina from four Bali bulls raised in Center Insemination Banjarbaru, province of south Kalimantan-Indonesia. All the bulls of at least 4-6 years of age and average weight of 625-650 kg were raised under a similar grazing system (various kinds of grass) and supplemented with concentrate at the rate of 4.5 kg/head/day. Two ejaculates were obtained from each bulls every week for one month. Immediately after semen collection, the ejaculates were immersed in the warm water bath at 37°C until their assessments in the laboratory. Then, semen parameters were assessed based on the macroscopic and microscopic characteristics. Macroscopic evaluations included volume, pH and color. Microscopic evaluations included live sperm rate, motility and abnormality. The main extender consisted of 10% skim milk, 5% egg-yolk, 1% glucose, 8% glycerol, Streptomycin (1 mg/mL) and Penicillin (1000 IU/mL).

This study was designed to compare the effect different concentrations of fish oil (liquid) with doses 0 (control), 50 mg, 100 mg, 150 mg and 200 mg in the 100 mL skim milk-egg yolk extender. The cryoprotective extender for the control group was the same as that for the treatment groups except that it was not supplemented with fish oil [(natural, 1000 mg) containing Omega-3 Marine; Triglycerides (300 mg) as EPA (180 mg) and DHA (120 mg) from Nature's Care (Manufacture Pty, Ltd. Minna Close Belrose, Australia)].

Freezing, Thawing and Evaluation

Semen was diluted to obtain a final concentration of 25×10^6 sperm/mL. Diluted semen was packed into 0.25 mL straws (Biovet, France). After sealing, straws were placed horizontally on a cold rack (5°C) and lowered into nitrogen vapors (-50°C), 4 cm above the surface of liquid nitrogen. After 3 min, when the temperature reached -120°C, the frozen straws were transferred into goblets of appropriate size and transferred into a liquid nitrogen tank. After two weeks of storage, the straws per treatment were randomly chosen, thawed in water at 37°C for 30 sec and evaluated for post-thawed semen quality including live sperm rate, motility and abnormality.

To evaluate sperm motility, a small droplet ($\approx 10 \mu\text{L}$) of post-thawed semen was placed in the center of a pre-warmed slide and covered with cover slip. It was transferred to a heated microscope stage at 37°C and subjectively calculated by phase contrast microscopy (400× magnification). For assessment of live sperm percentage, a modification of the eosin-nigrosin procedure described by Evans and Maxwell³⁰ was used. $10 \mu\text{L}$ of post-thawed semen was mixed with $30 \mu\text{L}$ of eosin-nigrosin stain. A thin smear was prepared and 200 spermatozoa was observed under a microscope (400 x). The same eosin-nigrosin stained slides were also used to define sperm abnormality for which 200 spermatozoa were examined for defects associated with tail region, mid-piece and sperm head.

Statistical Analysis

The data are expressed as mean \pm SEM. A one-way analysis of variance (ANOVA) with Tukey's multiple comparison test was applied to determine differences among the treatments using SPSS statistical software (version 16.0). Differences with P value 5% were considered to be statistically significant.


RESULTS

The effect of addition of different doses of fish oil on post-thawed live sperm rate, motility and abnormality is shown in Table 1 and Fig. 1. Addition of 100 mg fish oil to the 100 mL extender significantly ($P < 0.05$) increased both live sperm rate and motility when compared to the control and other treatments. However, the treatment with 200 mg fish oil caused significant ($P < 0.05$) reduction in live sperm

Table 1. Post-thawed sperm parameters including different dose fish oil Bali bulls

Treatments	Sperm parameters		
	Live Sperm Rate (%)	Motility (%)	Abnormality (%)
Control	54.64±2.41 ^b	42.50±2.50 ^{ab}	18.83±23.06 ^a
50 mg/100 mL	57.32±2.15 ^{bc}	43.33±1.44 ^b	21.17±50.49 ^{ab}
100 mg/100 mL	60.36±1.76 ^c	46.67±1.42 ^c	23.66±17.01 ^b
150 mg/100 mL	53.34±2.25 ^b	43.33±1.37 ^b	22.67±48.05 ^b
200 mg/100 mL	47.30±2.02 ^a	40.01±0.21 ^a	22.33±09.21 ^b

^{a,b,c} Values in the same column with different superscripts indicate significant difference at $P < 0.05$, (n=24)

**Fig 1.** Characteristic fresh semen of Bali bulls spermatozoa (Mean±SEM) (n=24)

rate in comparison with the control and other treatments. In addition, 200 mg fish oil supplementation also caused significant ($P < 0.05$) reduction when compared with the other treatment groups but not control group. A significant ($P < 0.05$) increment was observed in abnormality rate of frozen-thawed spermatozoa in comparison with the control group when all doses of fish oil except 50 mg/100 mL were added.

DISCUSSION

In the present study was evaluated the effects of fish oil supplementation on post-thaw semen quality in Bali bull spermatozoa. We hypothesized that fish oil supplement can enhanced semen quality in terms of live sperm rate and motility but not on sperm abnormality. The results obtained from this study are generally in agreement with the findings of previous studies conducted in other species such as turkeys [10], pigs [8] and rams [10], in which fish oil supplement improved semen quality. Meanwhile, reactive oxygen species (ROS) produced from cryopreservation

can also induce acrosome reaction and spermatozoa damage [19]. The toxicity effects of ROS eventually results in protein ionization and inactivation, peroxidation of lipids, in particular polyunsaturated fatty acids, and DNA damage which destabilize spermatozoa plasma membrane [20,21].

Live sperm rate and motility of semen are important factors because spermatozoa must travel from the vagina and uterus if semen is deposited by natural mating and AI, respectively. Cryopreservation, freezing and thawing processes may induce spermatozoa damages especially to the plasma membrane and organelles [22,23]. The determination of live sperm rate and motility is important for the evaluation of post-thaw quality of semen. In the present study, it was observed that addition of 100 mg fish oil to the 100 mL extender significantly increased both live sperm rate and motility, but treatment with 200 mg fish oil caused significant reduction in live sperm rate in comparison with the control and other treatments. In addition, 200 mg fish oil supplementation also caused significant ($P < 0.05$) reduction when compared with the other treatment groups but not control group. This situation is shown that the dose of 100 mg fish oil provides an ideal concentration in diluent so as to provide comfort to the spermatozoa, thus preventing damage to the process freezing, cryopreservation and thawing. These results are confirmed by previous reports by Abdi-Benemar et al. [24], who revealed that all frozen-thawed sperm characteristics are significantly enhanced with a surge in DHA levels from fish oil. One of the components of fish oil is PUFA with long chains and has been found in the spermatozoa of various species including ram, man and bull. These fatty acids improve the fluidity of the sperm plasma membrane which is then responsible for increased resistance of the sperm to cold conditions [25]. On the other hand, DHA can be presented as a cryoprotectant that will be able to penetrate the plasma membrane of the sperm and thus act intracellularly. Therefore, modifications in fatty acid composition of surrounding diluents with fish oil supplement can cause changes in the fluidity and elasticity of the sperm plasma membrane by incorporation its fatty acids in the lipids of sperm [26]. In the present study, 150 and 200 mg fish oil supplementations were observed to decrease the percentages of live sperm and motility. This result are

supported by the data previously reported by Kandelousi et al.^[27], who stated that higher levels of omega-6 fatty acids result in decreased sperm concentration, motility, and altered morphology.

In the present study, a significant increment was observed in abnormality rate of frozen-thawed spermatozoa in comparison with the control group when all doses of fish oil except 50 mg/100 mL were added. Majority of sperm abnormalities detected in this research were minor or secondary defects like simple coiled or bent tails that usually encounter in response to change in temperature, but not the defects of major type such as acrosome defect, defects associated through the sperm head and mid-piece, which are irreversible and genetically related^[28].

In conclusion, the results indicate that addition of 100 mg fish oil to 240 mL of skim milk-egg yolk extender could be beneficial for the improvement of the post-thaw quality of Bali bull spermatozoa including live sperm and motility. Therefore, fish oil addition at the dose of 100 mg per 100 mL skim milk-egg yolk extender can be recommended to bovine semen production centers.

REFERENCES

- Demir K, Öztürk Bakırer G, Cirit Ü, Bozkurt HH, Aktaş A, Bırlar S, Ak K, Pabuçcuoğlu S: Effects of cooling rate on membrane integrity and motility parameters of cryopreserved ram spermatozoa. *Kafkas Univ Vet Fak Derg*, 21 (1): 61-67, 2015. DOI: 10.9775/kvfd.2014.11726
- Isachenko E: Vitrification of mammalian spermatozoa in the absence of cryoprotectants: From past practical difficulties to present. *Reprod Biol*, 6, 191-200, 2003. DOI: 10.1016/S1472-6483(10)61710-5
- Khalili B, Farshad A, Zamiri MJ, Rashidi A, Fazeli P: Effects of sucrose and trehalose on the freezability of Markhoz goat spermatozoa. *Asian Australas J Anim Sci*, 22, 1614-1619, 2009. DOI: 10.5713/ajas.2009.90286
- Purdy PH: A review on goat sperm cryopreservation. *Small Ruminant Res*, 63, 215-225, 2006. DOI: 10.1016/j.smallrumres.2005.02.015
- Hammerstedt RH: Maintenance of bioenergetic balance in sperm and prevention of lipid peroxidation: A review of the effect on design of storage preservation systems. *Reprod Fertil Dev*, 5, 675-690, 1993. DOI: 10.1071/RD9930675
- Mandal R, Damodar B, Jitamanyu C: Role of membrane lipid fatty acids in sperm cryopreservation. *Adv Androl*, 2014 (2014), Article ID: 190542. DOI: 10.1155/2014/190542
- Stoeckel K, Nielsen LH, Fuhrmann H, Bachmann L: Fatty acid patterns of dog erythrocyte membranes after feeding of a fish-oil based DHA-rich supplement with a base diet low in n-3 fatty acids versus a diet containing added n-3 fatty acids. *Acta Vet Scand*, 6, 53-57, 2011. DOI: 10.1186/1751-0147-53-57
- Nichols PD, Glencross B, Petrie JR, Singh SP: Readily available sources of long-chain omega-3 oils: Is farmed Australian seafood a better source of the good oil than wild-caught seafood. *Nutrients*, 6, 1063-1079, 2014. DOI: 10.3390/nu6031063
- Rooke JA, Shao CC, Speake BK: Effects of feeding tuna oil on the lipid composition of pig spermatozoa and *in vitro* characteristics of semen. *Reproduction*, 121 (2): 315-322, 2001.
- Blesbois E, Douard V, Germain M, Boniface P, Pellet F: Effects of n-3 polyunsaturated dietary supplementation on the reproductive capacity of male turkeys. *Theriogenology*, 61, 537-549, 2004. DOI: 10.1016/S0093-691X(03)00207-3
- Samadian F, Towhidi A, Rezayazdi K, Bahreini M: Effects of dietary n-3 fatty acids on characteristics and lipid composition of ovine sperm. *Animal*, 4 (12): 2017-2022, 2010. DOI: 10.1017/S1751731110001308
- Castellano CA, Audet I, Laforest JP, Matte JJ, Suh M: Fish oil diets alter the phospholipid balance, fatty acid composition, and steroid hormone concentrations in testes of adult pigs. *Theriogenology*, 76, 1134-1145, 2011. DOI: 10.1016/j.theriogenology.2011.05.022
- Aksoy Y, Aksoy H, Altinkaynak K, Aydin HR, Ozkan A: Sperm fatty acid composition in subfertile men. *Prostaglandins Leukot Essent Fatty Acids*, 75, 75-79, 2006. DOI: 10.1016/j.plefa.2006.06.002
- Am-in N, Kirkwood RN, Techakumphu M, Tantasuparuk W: Lipid profiles of sperm and seminal plasma from boars having normal or low sperm motility. *Theriogenology*, 75, 897-903, 2011. DOI: 10.1016/j.theriogenology.2010.10.032
- Kaeoket K, Sang-urai P, Thamniyom A, Chanapiwat P, Techakumphu M: Effect of docosahexaenoic acid on quality of cryopreserved boar semen in different breeds. *Reprod Domest Anim*, 45, 458-463, 2010. DOI: 10.1111/j.1439-0531.2008.01239.x
- Maldjian A, Pizzi F, Gliozzi T, Cerolini S, Penny P, Noble R: Changes in sperm quality and lipid composition during cryopreservation of boar semen. *Theriogenology*, 63, 411-421, 2005. DOI: 10.1016/j.theriogenology.2004.09.021
- Chanapiwat P, Kaeoket K, Tummaruk P: Effects of DHA-enriched hen egg yolk and L-cysteine supplementation on quality of cryopreserved boar semen. *Asian J Androl*, 11, 600-608, 2009. DOI: 10.1038/aja.2009.40
- Evans G, Maxwell WMS: Salomon's Artificial Insemination of Sheep and Goats. University Press, Sydney, NSW, Australia, 1987.
- Mammoto A, Masumoto N, Tahara M, Ikebuchi Y, Ohmichi M, Tasaka K, Miyake A: Reactive oxygen species block sperm-egg fusion via oxidation of sperm sulfhydryl proteins in mice. *Biol Reprod*, 55, 1063-1068, 1996.
- Aitken RJ: Free radicals, lipid peroxidation and sperm function. *Reprod Fertil Dev*, 7, 659-668, 1995. DOI: 10.1071/RD9950659
- Aitken RJ, Gordon E, Harkiss D, Twigg JP, Milne P, Jennings Z, Irvine DS: Relative impact of oxidative stress on the functional competence and genomic integrity of humanspermatozoa. *Biol Reprod*, 59, 1037-1046, 1998. DOI: 10.1095/biolreprod59.5.1037
- Esteso MC, Fernández SMR, Soler AJ, Garde JJ: Head dimensions of cryopreserved red deer spermatozoa are affected by the thawing procedure. *Cryoletters*, 24, 261-268, 2003.
- Munoz OVL, Amirat-Briand T, Diaz L, Va'squez E, Schmidt S, Desherces M, Anton D, Bencharif D, Tainturier D: Effect of semen dilution to low-sperm number per dose on motility and functionality of cryopreserved bovine spermatozoa using low-density lipoproteins (LDL) extender: Comparison to Triladyl® and Bioxcell®. *Theriogenology*, 71, 895-900, 2009. DOI: 10.1016/j.theriogenology.2008.10.010
- Abdi-Benemar H, M Jafaroghli B, Khalili, MJ Zamiri, H. Ezazi, AA Shadparvar: Effects of DHA supplementation of the extender containing egg yolk and - tocopherol on the freezability and post-thawing fertility of ram semen. *Small Ruminant Res*, 130, 166-170, 2015. DOI: 10.1016/j.smallrumres.2015.06.013
- Wathes DC, Abayasekara DR, Aitken RJ: Polyunsaturated fatty acids in male and female reproduction. *Biol Reprod*, 77, 190-201, 2007. DOI: 10.1095/biolreprod.107.060558
- Conquer JA, Martin JB, Tummon I, Watson L, Tekpetey F: Effect of DHA supplementation on DHA status and sperm motility in asthenozoospermic males. *Lipids*, 35, 149-154, 2000. DOI: 10.1007/BF02664764
- Kandelousi SMA, Arshami J, Naserian AA, Abavisani A: The effects of addition of omega-3, 6, 9 fatty acids on the quality of bovine chilled and frozen-thawed sperm. *Open Vet*, 3 (1): 47-52, 2013.
- Chenoweth PJ: Genetic sperm defects. *Theriogenology*, 64, 457-468, 2005. DOI: 10.1016/j.theriogenology.2005.05.005

Effect of Fish Oil Addition to the Skim Milk-egg Yolk Extender on the Quality of Frozen-thawed Bali Bull Spermatozoa

ORIGINALITY REPORT

20%
SIMILARITY INDEX

12%
INTERNET SOURCES

18%
PUBLICATIONS

4%
STUDENT PAPERS

PRIMARY SOURCES

- 1** OĞUZ KARAKAŞ, Fatma, OĞUZ, Mustafa Numan and BÜYÜKOĞLU, Tülay. "The effects of using chicken drippings oil instead of the sunflower oil on performance, blood parameters, cholesterol and fatty acid composition of egg yolk in laying japanese quail (*Coturnix coturnix japonica*)", Kafkas Üniversitesi, 2012.
Publication **1%**
- 2** www.e-sciencecentral.org
Internet Source **1%**
- 3** www.theses.fr
Internet Source **1%**
- 4** pubmed.ncbi.nlm.nih.gov
Internet Source **1%**
- 5** Ghadeer Sabah Bustani, Falah Hasan Baiee. "Semen extenders: An evaluative overview of preservative mechanisms of semen and semen extenders", Veterinary World, 2021
Publication **1%**

6

Panida Chanapiwat. "Effects of DHA-enriched hen egg yolk and L-cysteine supplementation on quality of cryopreserved boar semen", *Asian Journal of Andrology*, 07/27/2009

Publication

1 %

7

Samaneh Ghasemi Fard, Fenglei Wang, Andrew J. Sinclair, Glenn Elliott, Giovanni M. Turchini. "How does high DHA fish oil affect health? A systematic review of evidence", *Critical Reviews in Food Science and Nutrition*, 2018

Publication

1 %

8

TAA Khalifa. "Testing Usability of Butylated Hydroxytoluene in Conservation of Goat Semen", *Reproduction in Domestic Animals*, 2/19/2008

Publication

1 %

9

Submitted to University of Limerick

Student Paper

1 %

10

www.thaiscience.info

Internet Source

1 %

11

Ijaz, A.. "Butylated hydroxytoluene inclusion in semen extender improves the post-thawed semen quality of Nili-Ravi buffalo (*Bubalus bubalis*)", *Theriogenology*, 200905

Publication

1 %

12

www.spencer-supplies.co.uk

Internet Source

1 %

13

www.gssrr.org

Internet Source

<1 %

14

journals.rifst.ac.ir

Internet Source

<1 %

15

research.dvs.gov.my

Internet Source

<1 %

16

www.science.gov

Internet Source

<1 %

17

A. Rastegarnia, A. Shahverdi, T. Rezaei topraggaleah, V. Shafiepour. "In vitro comparison of soybean lecithin-based extenders for cryopreservation of buffalo (*Bubalus bubalis*) semen", *Comparative Clinical Pathology*, 2013

Publication

<1 %

18

Kiernan, M., A. G. Fahey, and S. Fair. "The effect of the in vitro supplementation of exogenous long-chain fatty acids on bovine sperm cell function", *Reproduction Fertility and Development*, 2012.

Publication

<1 %

19

digital.car.chula.ac.th

Internet Source

<1 %

repositorio.ufla.br

21

Ali Khoshvaght, Armin Towhidi, Ahmad Zare-shahneh, Mohammad Noruozi, Mahdi Zhandi, Navid Dadashpour Davachi, Reza Karimi. "Dietary n-3 PUFAs improve fresh and post-thaw semen quality in Holstein bulls via alteration of sperm fatty acid composition", *Theriogenology*, 2016

Publication

<1 %

22

Chongshan Yuan, Hongtao Wang, Xichun Li, Hongyu Liu, Jing Zhao, Wenfa Lu, Jun Wang. "Combined Effect of Flaxseed Oil and Vitamin E Supplementation During Bull Semen Cryopreservation on Sperm Characteristics", *Biopreservation and Biobanking*, 2022

Publication

<1 %

23

Gliozzi, T.M.. "Quality and lipid composition of spermatozoa in rabbits fed DHA and vitamin E rich diets", *Theriogenology*, 20090401

Publication

<1 %

24

Submitted to Higher Education Commission Pakistan

Student Paper

<1 %

25

J Rubio-Guillén. "Effects of Cryopreservation on Bull Spermatozoa Distribution in

<1 %

Morphometrically Distinct Subpopulations", Reproduction in Domestic Animals, 8/2007

Publication

26

Malo, C.. "Antioxidant effect of rosemary (Rosmarinus officinalis) on boar epididymal spermatozoa during cryopreservation", Theriogenology, 201106

Publication

<1 %

27

ruidera.uclm.es

Internet Source

<1 %

28

www.agriculturejournals.cz

Internet Source

<1 %

29

Bedoor A. Salim, Muhsin S. G. AL-Moziel, Ausama Ayob Jaccob. "Sub-chronic effects of mefenamic acid alone or in combination with diclofenac on the female reproductive system in albino rats", Toxicology and Environmental Health Sciences, 2022

Publication

<1 %

30

Faraji, Z.. "Effect of feeding olive-pulp silage on the seminal characteristics and fatty acid profile of the sperm plasma membrane in Iranian fat-tailed sheep", Small Ruminant Research, 201206

Publication

<1 %

31

J.-H. Hu, L.-S. Zan, X.-L. Zhao, Q.-W. Li, Z.-L. Jiang, Y.-K. Li, X. Li. "Effects of trehalose

<1 %

supplementation on semen quality and oxidative stress variables in frozen-thawed bovine semen1", Journal of Animal Science, 2010

Publication

32

Jafaroghli, M.. "The effect of supplementation of cryopreservation diluents with sugars on the post-thawing fertility of ram semen", Small Ruminant Research, 201103

Publication

33

Kumaresan, A.. "Preservation of boar semen at 18°C induces lipid peroxidation and apoptosis like changes in spermatozoa", Animal Reproduction Science, 200901

Publication

34

Macaluso, Filippo, Rosario Barone, Patrizia Catanese, Francesco Carini, Luigi Rizzuto, Felicia Farina, and Valentina Di Felice. "Do Fat Supplements Increase Physical Performance?", Nutrients, 2013.

Publication

35

Masoudi, R., M. Sharafi, A. Zare Shahneh, A. Towhidi, H. Kohram, M. Zhandi, V. Esmaili, and A. Shahverdi. "Effect of dietary fish oil supplementation on ram semen freeze ability and fertility using soybean lecithin- and egg yolk-based extenders", Theriogenology, 2016.

Publication

<1 %

<1 %

<1 %

<1 %

36

Mohamed Elsayy, Tarek Sadaka, Mohamed Shourrap. "EFFECT OF UNSATURATED FATTY ACIDS ON HORMONAL PROFILE, ANTIOXIDANT STATUS, SEMEN QUALITY AND REPRODUCTIVE PERFORMANCE IN AGING BUCK RABBITS", Egyptian Journal of Rabbit Science, 2018

Publication

<1 %

37

Monika Trzcińska, Magdalena Baryła. "The use of butylated hydroxytoluene in cryopreservation of boar semen", Roczniki Naukowe Polskiego Towarzystwa Zootechnicznego, 2016

Publication

<1 %

38

Vinod Kumar Yata. "Chapter 6 Semen Extenders for Preservation of Sorted Semen", Springer Science and Business Media LLC, 2022

Publication

<1 %

39

liu.diva-portal.org

Internet Source

<1 %

40

pdfs.semanticscholar.org

Internet Source

<1 %

41

rep.bntu.by

Internet Source

<1 %

42

tesi.cab.unipd.it

Internet Source

<1 %

43

B K D BORAH, B C DEKA, R K BISWAS, P CHAKRAVARTY, S SINHA, K AHMED, S DEORI. "Antioxidants improve the semen quality following cryopreservation in Indian yak bulls", The Indian Journal of Animal Sciences, 2021

Publication

<1 %

44

Bligh, E. G., and W. J. Dyer. "A RAPID METHOD OF TOTAL LIPID EXTRACTION AND PURIFICATION", Canadian Journal of Biochemistry and Physiology, 1959.

Publication

<1 %

45

Camille Gautier, Christine Aurich. "'Fine feathers make fine birds" – The mammalian sperm plasma membrane lipid composition and effects on assisted reproduction", Animal Reproduction Science, 2022

Publication

<1 %

46

Dina M. Shokry, Mabrouk Attia Abd Eldaim, Magdy R. Badr, Hanem K. Khalifa, Sahar H. Orabi, Aziza M. Hassan, Ragab Dohreig. " Enhancement impact of leaves extract–base extender on cryopreservation and fertilization of Barki ram sperms: comparative study with vitamin E and selenium combination ", Italian Journal of Animal Science, 2021

Publication

<1 %

47

Larbi Allai, Anass Benmoula, Maia Marciane da Silva, Boubker Nasser, Bouchra El Amiri. "Supplementation of ram semen extender to improve seminal quality and fertility rate", Animal Reproduction Science, 2018

Publication

<1 %

48

Morteza Zarei, Behnam Rostami, Reza Masoumi, Mohsen Sharafi, Mohammad Hossein Shahir, Michael Stear. "Does addition of enriched egg yolks with different dietary oils to Tris-based extender improve ram sperm vital parameters in 5days liquid storage of semen?", Small Ruminant Research, 2018

Publication

<1 %

49

agrimaroc.org

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On