

# Increased Productivity of Eggs, FCR, and IOFC from Alabio Ducks through Addition of Artificial Light Ahemeral Method Using the Combination of the Intensity and Color of Monochromatic LED Light

*by* Tintin Rostini

---

**Submission date:** 07-Apr-2020 01:09PM (UTC+0700)

**Submission ID:** 1291722242

**File name:** IJB-Vol-15-No-5-p-155-162.pdf (451.06K)

**Word count:** 4555

**Character count:** 22296



INNSPUB

## RESEARCH PAPER

## OPEN ACCESS

## Increased Productivity of Eggs, FCR, and IOFC from Alabio Ducks through Addition of Artificial Light Ahemeral Method Using the Combination of the Intensity and Color of Monochromatic LED Light

Danang Biyatmoko<sup>1\*</sup>, Anis Wahdi<sup>1</sup>, Tintin Rostini<sup>2</sup>

<sup>1</sup>Department of Animal Science, Lambung Mangkurat University, Banjarbaru, Indonesia

<sup>2</sup>Department of Animal Science, Islamic University of Kalimantan Banjarmasin, Indonesia

**Key words:** Intensity, Light color, Ahemeral method, Duck productivity, LED.

<http://dx.doi.org/10.12692/ijb/15.5.155-162>

Article published on November 15, 2019

### Abstract

Light illumination, both color and intensity is one of the environmental factors that play an important role in regulating animal biorhythms. The use of artificial light in poultry produces very diverse responses, especially in metabolism, reproduction, and behavior. The purpose of this study was to determine the increase in egg productivity, feed conversion ratio (FCR) and income over feed cost (IOFC) of Alabio ducks through the addition of artificial light using the ahemeral method through a combination of intensity and color of monochromatic LED light. The research method uses a completely randomized factorial design of 4×3 with 3 replications. The first factor is the color of light (W) consisting of 4 levels including W1 (yellow), W2 (red), W3 (blue), W4 (green), and the second factor is light intensity (T) consisting of 3 levels including T1 (10 lux), T2 (15 lux) and T3 (20 lux). Nutritional content of crude protein (CP) 18%, metabolic energy (ME) 2800 kcal. kg<sup>-1</sup>. The variables observed were egg production, total egg weight, feed consumption, FCR and IOFC. The length of study and measurement of duck egg production is 4 weeks. The results showed that in the 4 weeks of observation the best treatment was a combination of blue light treatment with a light intensity of 15 lux (W3T2), with the highest egg production achievement of 22.40 ± 4.5 eggs.duck<sup>-1</sup>, the total egg weight achievement was 1373, 12 ± 16.5 g.duck<sup>-1</sup>, feed consumption of 810.12 ± 13.5 g.duck<sup>-1</sup>, FCR value of 4.13 ± 1.3 and IOFC of 29,405.39 ± 333, 1 Rp.duck<sup>-1</sup>. The conclusion of this study is the color of blue light with an intensity of 15 lux (W3T2) can increase the productivity of Alabio duck eggs and IOFC obtained.

\*Corresponding Author: Danang Biyatmoko ✉ [danangbiyatmoko@ulm.ac.id](mailto:danangbiyatmoko@ulm.ac.id)

## Introduction

The position of Alabio ducks is very important for the people in South Kalimantan because of their dependence on the egg products from the ducks. The contribution of eggs from Alabio ducks contributes more than 50% to the entire production of poultry eggs. As a cheap source of animal protein, demand for products from Alabio ducks continues to increase and even cross to other provinces in Indonesia, even to neighboring countries such as Malaysia, Brunei and not even rumored to Japan. Increased productivity of alabio duck eggs should be pursued so that the current total population of around 4.23 million in 2018 can meet high demand (BPS South Kalimantan, 2019). Genetic improvement has been carried out since 2010 by offices and centers, so other technological touches need to be adopted to be more productive and profitable (Biyatmoko, 2016).

One effort that can be done is through the adoption of artificial light, through the addition of light at night to increase the length of the day. Light is important because it is directly related to egg production. Color and light intensity are the most critical factors of all environmental factors because they are able to stimulate several gonadotropin hormones that control reproduction. Light intensity, light color and photoperiodicity (irradiation length) have important effects on egg production activities (Liu *et al.*, 2015). The hypothalamus will develop with a positive trigger of light stimuli. The light will emit electromagnetic waves which stimulate the ducks to bluff hypothalamic activity. Color is defined as the wavelength of light that stimulates the retina of the eye to produce changes in poultry behavior such as activity, aggressiveness and eating behavior (Svobodova *et al.*, 2015). Meanwhile, sufficient light intensity is required for the occurrence of releasing reproductive hormones such as FSH, LH as well as the synthesis and secretion of estradiol so as to guarantee the number of functional (mature) ovarian follicles and guarantee sustainable egg production (Liu *et al.*, 2015). According to Shabiha *et al.* (2013) and Svobodova *et al.* (2015), intensities that support poultry egg production range from 10 - 20 lux. To

increase production requires a certain wavelength so the wavelength of each color of light that is given will determine the response to the production of certain poultry eggs (Olanrewaju *et al.*, 2012).

Monochromatic light is a type of visible light with a single wavelength frequency and the distance between wavelengths is not too large. The electromagnetic spectrum of monochromatic light has a single color based on its wavelength, namely red (630-760 nm), orange (590-630 nm), yellow (570-590 nm), green (500-570 nm), blue (450-70 nm) 500 nm, and purple (400-450 nm) (Elert, 2008). Kasiyati (2018) <sup>4</sup> said that the successful use of poultry irradiation with light emitting diode (LED) lamps as a source of light is mostly done on modern farms, because it provides advantages in electrical energy efficiency, color emission is more stable, brighter, longer life (long life), and can reduce the humidity of the cage. Artificial light is not only a source of illumination in modern poultry farming but also plays a role in the reproduction response and production of poultry. The study of several colors of light is important because it is possible to cause different responses from production variables.

<sup>5</sup> The purpose of this study was to determine the increase in egg production, total egg weight, feed consumption, FCR and income over feed cost (IOFC) Alabio Ducks through the addition of artificial light using the ahemeral method through a combination of intensity and color of monochromatic LED light.

## Material and methods

The study was carried out in poultry cages where measurements of egg production and research observations were carried out for four weeks.

The research material uses Alabio Ducks so that there are 180 ducks, using laying ducks aged 7 months after entering the egg production cycle. The nutritional content of the experimental rations prepared iso protein and iso energy including crude protein (CP) 18%, Metabolizable Energy (ME) 2800 kcal. kg<sup>-1</sup>, SK 6%, Calcium 2.5%, and Phosphorus (P) 1.5% using the



Win Feed program version 2.8. Drinking water is given to ducks on an ad libitum basis.

The experimental design used a completely randomized  $4 \times 3$  factorial pattern design with 3 replications which consisted of 5 layer ducks for each repetition. The research treatment consisted of 2 factors, where the first factor was the color of the light (W) consisting of 4 levels namely W1 (yellow), W2 (red), W3 (blue), W4 (green), and the second factor namely Light intensity (T) consisting of 3 the standards are T1 (10 lux), T2 (15 lux) and T3 (20 lux). Additional light is given six hours with day length of 28 hours consisting of daylight 18 hours of light and 10 hours of dark is an application with the ahemeral method. Variables observed were egg production,

total egg weight (g.duck<sup>-1</sup>), feed consumption (g.duck<sup>-1</sup>), Feed Conversion Ratio (FCR), and Income Over Feed Cost (IOFC). Data obtained in this study were analyzed by variance analysis (Anova). If the analysis of variance is significantly different, it will be continued with the mean difference test using the Duncan Multiple Range Test (DMRT) at the 5% and 1% significance levels (Steel and Torrie, 1993).

### Results and discussion

The results of research observations on the effect of the addition of artificial light ahemeral method using a combination of intensity and color of monochromatic LED light on egg production, total egg weight, feed consumption, and FCR of Alabio ducks are presented in Table 1.

**Table 1.** Average egg production, feed consumption and feed conversion ratio (FCR) of Alabio Ducks for four weeks.

Treatment	Egg Production (egg.duck <sup>-1</sup> )	Total Egg Weight (g.duck <sup>-1</sup> )	Feed Consumption (g.duck <sup>-1</sup> )	FCR
W1T1 (yellow color, 10 lux)	13,06 ± 4,5 <sup>a</sup>	800,98 ± 11,6 <sup>a</sup>	736,88 ± 10,6 <sup>a</sup>	6,44 ± 1,1 <sup>a</sup>
W1T2 (yellow color, 15 lux)	13,06 ± 5,7 <sup>a</sup>	800,98 ± 13,7 <sup>a</sup>	727,72 ± 9,7 <sup>a</sup>	6,36 ± 1,3 <sup>a</sup>
W1T3 (yellow color, 20 lux)	14,93 ± 4,3 <sup>b</sup>	915,41 ± 14,3 <sup>b</sup>	802,92 ± 14,3 <sup>b</sup>	6,14 ± 1,1 <sup>b</sup>
W2T1 (red color, 10 lux)	16,80 ± 6,1 <sup>b</sup>	1029,84 ± 16,1 <sup>b</sup>	863,56 ± 12,1 <sup>b</sup>	5,87 ± 1,5 <sup>b</sup>
W2T2 (red color, 15 lux)	16,80 ± 5,4 <sup>b</sup>	1029,84 ± 15,4 <sup>b</sup>	857,68 ± 12,4 <sup>b</sup>	5,83 ± 2,2 <sup>b</sup>
W2T3 (red color, 20 lux)	18,66 ± 4,6 <sup>c</sup>	1144,26 ± 17,6 <sup>c</sup>	871,24 ± 15,6 <sup>c</sup>	5,33 ± 1,7 <sup>c</sup>
W3T1 (blue color, 10 lux)	18,66 ± 4,2 <sup>c</sup>	1144,26 ± 15,2 <sup>c</sup>	874,52 ± 12,2 <sup>c</sup>	5,35 ± 1,6 <sup>c</sup>
W3T2 (blue color, 15 lux)	22,40 ± 4,5 <sup>c</sup>	1373,12 ± 16,5 <sup>c</sup>	810,12 ± 13,5 <sup>c</sup>	4,13 ± 1,3 <sup>c</sup>
W3T3 (blue color, 20 lux)	20,53 ± 3,9 <sup>d</sup>	1258,69 ± 12,9 <sup>d</sup>	830,72 ± 11,9 <sup>d</sup>	4,62 ± 1,2 <sup>d</sup>
W4T1 (green color, 10 lux)	18,66 ± 4,9 <sup>c</sup>	1144,26 ± 14,9 <sup>c</sup>	885,96 ± 12,9 <sup>c</sup>	5,42 ± 1,6 <sup>c</sup>
W4T2 (green color, 15 lux)	20,53 ± 5,5 <sup>d</sup>	1258,69 ± 13,5 <sup>d</sup>	819,92 ± 12,5 <sup>d</sup>	4,56 ± 1,9 <sup>d</sup>
W4T3 (green color, 20 lux)	20,53 ± 6,1 <sup>d</sup>	1258,69 ± 14,1 <sup>d</sup>	827,12 ± 14,1 <sup>d</sup>	4,60 ± 1,9 <sup>d</sup>

Note: Figures followed by different superscripts in the same column show significantly different ( $p < 0.05$ ); W: Light color; T: Light intensity; FCR: Feed Conversion Ratio.

#### Egg Production and Total Weight

At the egg production level (egg) in Table 1, it was seen that during the four weeks of observation, the combination of light color with light intensity of monochromatic LED lamps with an ahemeral method significantly affected the amount of egg production in eggs ( $p < 0.05$ ), where treatment with the combination of blue light color and light intensity of 15 lux (W3T2) gave the highest results with the highest egg production achievement reaching  $22.40 \pm$

4.5 eggs.duck<sup>-1</sup> with the total egg weight achievement of  $1373.12 \pm 16.5$  g.duck<sup>-1</sup>, significant higher than other treatments with a production range of  $13.06 - 20.53$  eggs.ducks<sup>-1</sup> and weight achievements ranging from  $800.98 - 1258$  g.duck<sup>-1</sup> ( $p < 0.05$ ). The lowest productivity is produced by the combination of yellow light with a light intensity of 10 lux (W1T1) and 15 lux (W1T2) with egg production of  $13.06 \pm 4.6$  eggs.duck<sup>-1</sup>, respectively with a total egg weight of  $800, 98 \pm 11.6$  g.duck<sup>-1</sup>.

**Table 2.** The average business income or Income over Feed Cost (IOFC) of Alabio Ducks for four weeks.

Treatment	Salling Value of Eggs (Rp.duck <sup>-1</sup> )	Cost of Feed Consumption (Rp.ekor <sup>-1</sup> )	(IOFC) (Rp.duck <sup>-1</sup> )
W1T1 (yellow color, 10 lux)	32039,20 ± 311,6 <sup>a</sup>	23.212,40 ± 250,6 <sup>a</sup>	8.826,81 ± 121,1 <sup>a</sup>
W1T2 (yellow color, 15 lux)	32039,20 ± 312,7 <sup>a</sup>	22.924,03 ± 259,7 <sup>a</sup>	9.115,17 ± 115,3 <sup>a</sup>
W1T3 (yellow color, 20 lux)	36616,40 ± 334,3 <sup>b</sup>	25.292,74 ± 224,3 <sup>b</sup>	11.323,66 ± 161,1 <sup>b</sup>
W2T1 (red color, 10 lux)	41193,60 ± 476,1 <sup>b</sup>	27.203,20 ± 202,1 <sup>b</sup>	13.990,38 ± 189,5 <sup>b</sup>
W2T2 (red color, 15 lux)	41193,60 ± 415,4 <sup>b</sup>	27.017,82 ± 212,4 <sup>b</sup>	14.175,78 ± 170,2 <sup>b</sup>
W2T3 (red color, 20 lux)	45770,40 ± 427,6 <sup>c</sup>	27.445,05 ± 205,6 <sup>c</sup>	18.325,35 ± 211,7 <sup>c</sup>
W3T1 (blue color, 10 lux)	45770,40 ± 435,2 <sup>c</sup>	27.548,10 ± 312,2 <sup>c</sup>	18.222,35 ± 191,6 <sup>c</sup>
W3T2 (blue color, 15 lux)	54924,80 ± 446,5 <sup>e</sup>	25.519,41 ± 289,5 <sup>e</sup>	29.405,39 ± 333,1 <sup>e</sup>
W3T3 (blue color, 20 lux)	50347,60 ± 452,9 <sup>d</sup>	26.168,13 ± 261,9 <sup>d</sup>	24.179,47 ± 211,2 <sup>d</sup>
W4T1 (green color, 10 lux)	45770,40 ± 313,9 <sup>c</sup>	27.908,50 ± 212,9 <sup>c</sup>	17.861,94 ± 221,6 <sup>c</sup>
W4T2 (green color, 15 lux)	50347,60 ± 454,5 <sup>d</sup>	25.828,29 ± 212,5 <sup>d</sup>	24.519,31 ± 211,9 <sup>d</sup>
W4T3 (green color, 20 lux)	50347,60 ± 434,1 <sup>d</sup>	26.054,87 ± 214,1 <sup>d</sup>	24.292,74 ± 231,9 <sup>d</sup>

Note: Figures followed by different superscripts in the same column show significantly different ( $p < 0.05$ ), W: Light color; T: Light intensity; IOFC: Income Over Feed Cost; feed price of IDR 4500.kg<sup>-1</sup>; the price of duck egg Rp. 2500.egg<sup>-1</sup>; 1 kg of eggs is equivalent to 16 eggs of Alabio duck; one egg averaging 61,3 g.

This research shows that light that has short wavelengths such as blue light with wavelengths around 480 nm and green light with wavelengths of 520 nm are more effective at stimulating the retina of the eye to be able to stimulate the hypothalamus and pituitary gland in producing reproductive hormones namely the follicle stimulating hormone (FSH) and luteinizing hormone (LH) compared to the color of light with other long wavelengths such as yellow (580 nm) and red (700 nm) (Mohamed *et al.*, 2014), so that ovarian development and maturation is faster (Jin *et al.*, 2011; Hassan, *et al.*, 2013; Biyatmoko, 2014) While the best combination of light intensity of 15 lux achieved in producing egg production in this study is the recommended intensity range for laying poultry. Light at intensities of 10-20 lux is able to stimulate the process ovulation of ducks, while the color of light affects the wavelength it receives (Shabiha *et al.*, 2013). The difference in production levels due to the different colors and light intensities is in line with the results of research by Rozenboim *et al.* (2012) and Kim *et al.* (2012).

The color affects the behavior patterns of poultry, while sufficient light intensity will be able to bully the stimulation of the ovulation process of the egg that

initiates egg production. The lack of intensity will inhibit the stimulation of ovulation that occurs due to inhibition of the secretion of the gonadotropin hormone in poultry. Irradiating blue and green colors make poultry quieter, food use efficiency (FCR) better and can be fully converted into eggs (Rozenboim, *et al.*, 2012; Hassan, *et al.*, 2013). On the other hand, the yellow and red light colors that cause ducks are more aggressive and active so that the consumption of feed is widely used to help energy in poultry locomotion so that egg production decreases. Energy and protein efficiency is fully manifested in producing egg production so that in addition to higher production, the weight of eggs produced is also greater and heavier for ducks that get blue and green LED monochromatic light colors (Freitas *et al.*, 2010).

This result corroborates the statement of Kasiyati *et al.*, (2010) which states that in white light blue laying eggs with an intensity of 15-25 lux will increase serum estradiol concentration, followed by an increase in ovarian weight and follicular diameter F1, where the greater the ovary and follicle diameter which is formed then causes the yolk (vitelin) to be bigger and the greater the weight of the egg.



#### Feed consumption and feed conversion ratio (FCR)

In Table 1, it can be seen that the highest consumption of duck ration was produced by the treatment of green light with a light intensity of 10 lux (W4T1) of  $885.96 \pm 12.9$  g.duck<sup>-1</sup> with an average consumption of  $727.72 \pm 9.7$  g.duck<sup>-1</sup> ( $p < 0.05$ ). However, this consumption level indicator does not positively correlate with the level of duck egg production, because it is more determined by feed conversion (FCR) where the better the FCR (low FCR value), the higher and more efficient in egg production. Table 1 shows that the best FCR is produced by blue light treatment with a light intensity of 15 lux (W3T2) with an FCR value of  $4.13 \pm 1.3$  linear with the highest level of production, while the worst FCR is produced by yellow light treatment with intensity 10 lux light (W1T1) with FCR reaching  $6.44 \pm 1.1$  linear with the lowest egg production achievement of  $13.06 \pm 4.6$  g.duck<sup>-1</sup> is the same as W1T2 treatment egg production (yellow light color, intensity of 15 lux).

In general, the highest consumption of ration was produced by irradiation with red light (W2) with an average of 864.16 g.duck<sup>-1</sup> is higher than other colors of light, followed by green light (W4) by 844.33 g, the blue color (W3) is 864.14 g.duck<sup>-1</sup> and the last color is yellow light (W1) is 755.84 g.duck<sup>-1</sup>.

The results of this study as stated by Zhu *et al.* (2019), where the use of red light can increase chicken locomotion which can trigger chicken aggressiveness so it is recommended to use it in the starter maintenance phase (brooding) because poultry is more active not in the egg production phase (Olanrewaju *et al.*, 2012). But if the intensity of light given is excessive can disrupt the daily rhythm (diurnal) egg production. Good irradiation in ducks other than the animal's preferred light color, blue and green, must also be balanced with adequate light intensity (number of lumens of light per watt) that ducks must receive, so that most of their feed consumption will be used fully for the egg production process (Li *et al.*, 2014). The vision of birds has a different sensitivity to the color of light where the

color received by the retina of the eye will be able to distinguish colors with different levels of sensitivity according to differences in wavelength (Firouzi, *et al.*, 2014). Wavelength influences the behavior of poultry (Kim *et al.*, 2012). Feeding behavior of poultry, including ducks given irradiation in blue and green, will be calmer (Shabiha *et al.*, 2013).

Borille *et al.* (2013) stated that irradiating yellow light was not effectively given to laying poultry because it would show the lowest production yields and even reduce the level of egg production and efficient use of food (feed conversion ratio). Meanwhile, the illumination of blue light turns out to reduce locomotion (movement activity) and stand on ducks so that ducks become calmer (Huber *et al.*, 2013; Hassan *et al.*, 2014). Instead according to Firouzi *et al.* (2014), light functions to regulate daily rhythm and several important functions in the body such as body temperature and various metabolic stages associated with feeding and digestion. Feed consumption is more efficiently used to produce fully egg production because ducks do not waste a lot of energy for their locomotion in energy activities (Olanrewaju *et al.*, 2012). This makes improvements to the feed conversion as shown in Table 1 where the treatment of blue light with an intensity of 15 lux (W3T2) results in the best feed conversion (FCR) of 4.13 while the treatment of yellow light with an intensity of 10 lux (W1T1) produces the worst FCR is 6.44.

In Indonesia as a tropical country, the periodicity of solar lighting is only around 12 hours day<sup>-1</sup> while the light requirement for poultry to be able to produce is 14-16 hours with sufficient intensity to stimulate the process of ovulation of eggs so the addition through artificial lighting (artificial light) at night is necessary (Biyatmoko, 2012). The addition of irradiation through the addition of six hours of light per day using ahemeral method to 18 hours of light and 10 hours of dark (one cycle of 28 hours.day<sup>-1</sup>) is quite promising, because it is proven to be able to stimulate ovulation and increase the secretion of FSH hormone so that it bluffs activation. egg cells (ovaries) for

oviposition or laying eggs (Hassan *et al.*, 2013) and significantly increase the yield of egg production and FCR of Alabio ducks in this study. These results reinforce the statement of Kasiyati (2018) that the use of longer bright light with a photoperiod of 16 bright hours: 8 hours dark (16L: 8D) increases the synthesis and secretion of estradiol in local poultry so that in the long run it can increase the number of functional ovarian follicles and guarantee sustainable egg production.

#### Income Over Feed Cost (IOFC)

Operating income or income over feed cost (IOFC) for four weeks of egg observation is presented in Table 2. Income over feed cost (IOFC) is income received from livestock businesses or in this case Alabio ducks are run, by looking at the margin of difference between the selling value of eggs produced with the cost of feed consumption (feed costs). The higher IOFC management, the higher the income generated and the more profitable. In Table 2, the highest income or IOFC is produced by the combination of blue light with a light intensity of 15 lux (W3T2) with an IOFC achievement of  $29,405.39 \pm 333.1$  Rp.duck<sup>-1</sup> based on egg selling value of  $54,924.80 \pm 446.5$  Rp.duck<sup>-1</sup> and feed costs were  $25,519.41 \pm 289.5$  Rp.duck<sup>-1</sup>, while the lowest IOFC was produced by the combination of yellow light color with 10 lux light intensity of  $8,826.81 \pm 121.1$  Rp.duck<sup>-1</sup> based on the sale value of eggs of  $32,039.20 \pm 311.6$  Rp.duck<sup>-1</sup> and the feed costs of  $23,212.40 \pm 250.6$  Rp.duck<sup>-1</sup>.

#### Conclusion

The combination of blue light color with a light intensity of 15 lux (W3T2) can increase the ability of Alabio duck egg productivity with the highest egg production achievement of  $22.40 \pm 4.5$  eggs.duck<sup>-1</sup>, the total egg weight achievement of  $1373.12 \pm 16.5$  g.duck<sup>-1</sup>, feed consumption of  $810.12 \pm 13.5$  g.duck<sup>-1</sup>, FCR value of  $4.13 \pm 1.3$  and income over feed cost (IOFC) of  $29,405.39 \pm 333.1$  Rp.duck<sup>-1</sup>.

#### Acknowledgements

The researcher would like to express his gratitude and highest appreciation for this PTUPT research funding

from the Ministry of Research, Technology and Higher Education Ministry of Education in 2019. Hopefully the results of this study will be useful and can improve scientific knowledge and encourage technology transfer from TKT in this study.

#### References

- BPS Kalimantan Selatan.** 2019. Indeks Harga Konsumen/ Inflasi Kalimantan Selatan. <https://www.bi.go.id/id/publikasi/kajianekonomiregional/kalsel/Contents/Kajian%20Ekonomi>.
- Biyatmoko D.** 2014. Effect the combinations of light color and intensity of light to egg first laying and production egg of Alabio laying duck. *International Journal of Biosciences* **5**, 80-85.
- Biyatmoko D.** 2016. The effect of protease enzyme supplementation to productivity eggs of alabio duck. *International Journal of Biosciences* **8(2)**, 202-208.
- Borille R, Garcia RG, Royer AFB, Santana, Colet S, Naas IA, Caldara FR, Almeida Paz ICL, Rosa ES, Castilho VAR.** 2013. The use of light-emitting diodes (LED) in commercial layer production. *Revista Brasileira de Ciência Avícola* **15**, 135-140. <https://doi.org/10.1590/S1516-635X2013000200009>
- Firouzi S, Haghbin Nazarpak H, Habibi H, Jalali SS, Nabizadeh Y, Rezaee F, Ardali R, Marzban M.** 2014. Effects of color lights on performance, immune response and hematological indices of broilers. *Journal World's Poultry Research* **4**, 52-55.
- Freitas HJ, Cotta JTB, Oliveira AL, Murgas LDS, Gewehr CE.** 2010. Effects of different lighting programs in laying hens housed in free range. *Revista Biotemas* **23**, 157-162. (in Portuguese).
- Hassan MR, Sultana S, Choe HS, Ryu KS.** 2014. Effect of combinations of monochromatic led light color on the performance and behavior of laying hens. *Journal of Poultry Science* **51**, 321-326.



- Hassan RMd, Sultana S, Sung HH, Saon RK.** 2013. Effect of monochromatic and combined light colour on performance, blood parameters, ovarian morphology and reproductive hormones in laying hens. *Italian Journal of Animal Science* **12**, 1–6.
- Huber-Eicher B, Suter A, Spring-Stahli P.** 2013. Effects of colored light-emitting diode illumination on behavior and performance of laying hens. *Journal of Poultry Science* **92**, 869-873.  
<https://doi.org/10.3382/ps.2012-02679>
- Jin E, Jia I, Li J, Yang G, Wang Z, Cao J, Chen Y.** 2011. Effect of Monochromatic Light on Melatonin Secretion and Arylalkylamine N-Acetyltransferase mRNA Expression in The Retina and Pineal Gland of Broilers. *The anatomical record* **294**, 1233–1241.
- Kasiyati N, Kusumorini H, Maheshwari W, Manalu.** 2010. Fotostimulasi cahaya monokromatik untuk optimasi reproduksi dan karakteristik karkas puyuh (*Coturnix coturnix japonica*). *Prosiding Hewan dalam Rangka Swasembada Pangan Nasional*. Bogor, 6-7 Oktober, p 118-122.
- Kasiyati.** 2018. The role of light in birds life: the response of growth and reproduction. *Buletin Anatomi dan Fisiologi* **3(1)**, 116 – 125.
- Kim MJ, Hossian MS, Nazma A, Jae CN, Han TB, Hwan KK, Dong WK, Chae HC, Choi HC, Ok SS.** 2012. Effect of monochromatic light on sexual maturity, production performance and egg quality of laying hens. *Avian Biology Research* **5**, 69–74.
- Li D, Zhang L, Yang M, Yin H, Xu H, Trask JS, Smith DG, Zhang Z, Zhu Q.** 2014. The effect of monochromatic light-emitting diode light on reproductive traits of laying hens. *Journal Applied Poultry Research* **23**, 367–375.
- Liu L, Li D, Gilbert ER, Xiao Q, Zhao X, Wang Y, Yin H, Zhu Q.** 2015. Effect of monochromatic light on expression of estrogen receptor (ER) and progesterone receptor (PR) in ovarian follicles of chicken. *Plos One* **10(12)**, 1-14.
- Mohamed R A, Elthoth MM, El-Saidy NR.** 2014. Rearing broiler chickens under monochromatic blue light improve performance and reduce fear and stress during pre-slaughter handling and transportation. *Biology in Animal Husbandry* **30(3)**, 457-471.
- Morrill WBB, Barnabé JMC, Da Silva TPN, Pandorfi H, Gouveia-Neto AS, Wellington SS.** 2014. The effect of RGB monochromatic and polychromatic LED lighting on growth performance, behavior, and development of broilers. *Proceedings of Society of Photo-Optical Instrumentation Engineers*; San Francisco, CA, USA.  
<http://dx.doi.org/10.1117/12.2036602>
- Olanrewaju HA, Purswell PL, Collier SD, Branton SL.** 2012. Influence of photoperiod, light intensity, and their interaction on growth performance and carcass characteristics of broilers grown to heavy weights. *International Journal of Poultry Science* **11**, 739–746.
- Rozenboim I, Mobarkey N, Avital-CN, Kashash-HY, Heiblum R, Chaiseha Y, Halawani ME.** 2012. Photostimulation effects on reproductive activities of domestic birds. In: *The XXIV World's Poultry Congress*; Bahia, Brazil, p 1–8.
- Shabiha S, Hassan R, Choe HS, Ryu KS.** 2013. Impact of different monochromatic LED light colour and bird age on the behavioral output and fear response in ducks. *Italian Journal of Animal Science* **12**, 573 – 584.
- Steel RGD, Torrie JH.** 1993. *Prinsip dan Prosedur Statistika*. Edisi 2. Terjemahan Sumantri B. PT Gramedia Pustaka Utama. Jakarta.
- Svobodova J, Tumova E, Chodova D.** 2015. Effect of light colour on egg production. *Czech Journal of Animal Science* **60(12)**, 550-556.



Zhu HX, Hu MD, Guo BB, Qu XL, Lei MM, Chen R, Chen Z, Shi ZD. 2019. Effect and molecular regulatory mechanism of monochromatic light colors on the egg-laying performance of

Yangzhou geese. *Animal Reproduction Science* **204**, 131-139.

<http://dx.doi:10.1016/j.anireprosci.2019.03.015>

# Increased Productivity of Eggs, FCR, and IOFC from Alabio Ducks through Addition of Artificial Light Ahemeral Method Using the Combination of the Intensity and Color of Monochromatic LED Light

## ORIGINALITY REPORT

7%

SIMILARITY INDEX

5%

INTERNET SOURCES

2%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.innspub.net](http://www.innspub.net)

Internet Source

5%

2

Submitted to Georgia State University

Student Paper

1%

3

[www.mdpi.com](http://www.mdpi.com)

Internet Source

1%

4

Farid N.K. Soliman, Karim El-Sabrout. "Light wavelengths/colors: Future prospects for broiler behavior and production", Journal of Veterinary Behavior, 2019

Publication

1%

5

Zurmiati ., Wizna ., M. Hafil Abba, Maria Endo Mahat. "Effect of the Balance of Energy and Protein in Rations Given to Pitalah Ducks along with the Probiotic Bacillus amyloliquefaciens on the Live Weight, Percentage of Carcass, Percentage of Abdominal Fat and Income Over

1%

# Feed Cost", International Journal of Poultry Science, 2017

Publication

---

---

Exclude quotes      On

Exclude matches      < 1%

Exclude bibliography      On